



University of Guilan

journal homepage: <https://cse.guilan.ac.ir/>

## Extended Auxiliary Equation Method to the perturbed Gerdjikov-Ivanov Equation

Jamilu Sabi'u<sup>1</sup>, Hadi Rezazadeh<sup>2,\*</sup>, Mostafa Eslami<sup>3</sup>, Alper Korkmaz<sup>4</sup><sup>1</sup> Department of Mathematics, Northwest University, Kano, Nigeria<sup>2</sup> Faculty of Engineering Technology, Amol University of Special Modern Technologies, Amol, Iran<sup>3</sup> Department of Mathematics, Faculty of Mathematical Sciences, University of Mazandaran, Babolsar, Iran<sup>4</sup> Nord Staße 9, Weimar, Germany

### ARTICLE INFO

#### Article history:

Received 5 January 2023

Received in revised form 25 January 2023

Accepted 6 February 2023

Available online 6 February 2023

#### Keywords:

Perturbed Gerdjikov–Ivanov equation

Travelling wave solution

Extended auxiliary equation method

Complex wave solution.

### ABSTRACT

We apply utilized the extended form of the auxiliary equation method to obtain extensively reliable exact travelling wave solutions of perturbed Gerdjikov–Ivanov equation (GIE) that is widely used as a model in the field theory of quanta and non-linear optics. The method is based on a simple first order second degree ODE. The new form of the approach gives more solutions to the governing equation efficiently.

## 1. Introduction

Optical soliton perturbation is studied in both physical and experimental phenomena modelled in to linear and non-linear PDEs such as; non-linear optics, electro-magnetics, plasma and solid state physics, water wave propagation in shallow media, fluid problems, and more. However, different approaches have been proposed over the years to handle the analytical solution of non-linear partial differential equations (NPDEs), see [1–18].

The perturbed GIE with full non-linearity is given in [13] of the form

\*Corresponding author: [H.rezazadeh@ausmt.ac.ir](mailto:H.rezazadeh@ausmt.ac.ir) (Hadi Rezazadeh),

E-mail address

[Sabiujamilu@gmail.com](mailto:Sabiujamilu@gmail.com) (Jamilu Sabi'u),

[H.rezazadeh@ausmt.ac.ir](mailto:H.rezazadeh@ausmt.ac.ir) (Hadi Rezazadeh),

[Mostafa.Eslami@umz.ac.ir](mailto:Mostafa.Eslami@umz.ac.ir) (Mostafa Eslami),

[Alper.Korkmaz@gmail.com](mailto:Alper.Korkmaz@gmail.com) (Alper Korkmaz).

$$i w_t + a w_{xx} + b |w^4| + i c w^2 w_x^* = i(\alpha w_t + \lambda (w |w|^{2m})_x) + \mu ((|w|^{2m})_x w), \quad (1.1)$$

herein  $a$ ,  $b$  and  $c$  are constants with real values and  $m$  denotes all non-linearity effects,  $\alpha$  is the dispersion coefficient,  $\mu$  is another dispersion term with higher effect as  $\lambda$  is the coefficients of short pulses that steepen themselves.

It is remarkable to mention that, quite number of methods have been set for finding the solutions of GIE. Structure in Hamiltonian forms, algebro-geometric solutions approach, tangent expansion technique, solution in Wronskian-types, transformations similar to Darboux. For more details see [19-24]. In this paper, we will propose another scheme for the exact and soliton solutions of GIE with aid of extended form of the auxiliary equation method (EAEM) [25].

The remaining article is designed as follows. In the next, we described methodology of the our scheme using EAEM in [25]. Then, in Section 3, some solutions are set to the GIE using the aforementioned methodology. Eventually, a brief conclusion is given in the last section.

## 2. Description of the method:

Suppose that we have the following PDE

$$F(u, u_t, u_x, u, \dots) = 0, \quad (2.1)$$

where  $u(x, t) = u$  is the dependent variable,  $x$  and  $t$  are dependent variables,  $F$  is a polynomial-like in terms of  $u$  and its derivatives.

EAEM mainly consists of three steps:

**Step 1:** The first step is the reduction of the governing equation by a compatible wave transformation such as

$$u(x, t) = U(\zeta), \quad \zeta = x - vt, \quad (2.2)$$

where  $v$  is arbitrary constant. This transform can be changed to some complex-valued transforms in order to the governing equation into a non-linear ODE with respect to the variable  $\zeta$  of the form

$$G(U, U', U'', \dots), \quad (2.3)$$

where the prime represents classical derivative for the independent variable  $\zeta$ .

**Step 2.** Assume that the predicted solution to Eq. (2.3) is of the form of finite series as

$$U(\zeta) = \sum_{i=0}^N n_i \Phi^i(\zeta) \quad (2.4)$$

in which  $n_i$  ( $i = 0, 1, 2, 3, \dots, N$ ) are all real constants to be determined,  $N$  is a positive integer which can be determined by classical balancing procedure between the highest order derivative terms and the highest degree non-linear terms in Eq. (2.3). One should note that  $\Phi(\zeta)$  satisfies the following new first order second degree auxiliary ODE

$$\left( \frac{d\Phi}{d\zeta} \right)^2 = m_1 \Phi^2(\zeta) + m_2 \Phi^4(\zeta) + m_3 \Phi^6(\zeta) \quad (2.5)$$

where  $m_1, m_2, m_3$  are real parameters to be found.

Equation (2.5) gives the following solutions:

When  $m_1 > 0$ ,

$$\Phi_1(\zeta) = \left( \frac{-m_1 m_2 \operatorname{sech}^2(\sqrt{m_1} \zeta)}{m_2^2 - m_1 m_3 (1 + \varepsilon \tanh(\sqrt{m_1} \zeta))^2} \right)^{\frac{1}{2}}. \quad (2.6)$$

When  $m_1 > 0$ ,

$$\Phi_2(\zeta) = \left( \frac{m_1 m_2 \operatorname{csch}^2(\sqrt{m_1} \zeta)}{m_2^2 - m_1 m_3 (1 + \varepsilon \coth(\sqrt{m_1} \zeta))^2} \right)^{\frac{1}{2}}. \quad (2.7)$$

When  $m_1 > 0, \Delta > 0$ ,

$$\Phi_3(\zeta) = \left( \frac{2m_1}{\varepsilon \sqrt{\Delta} \cosh(2\sqrt{m_1} \zeta) - m_2} \right)^{\frac{1}{2}}. \quad (2.8)$$

When  $m_1 < 0, \Delta > 0$ ,

$$\Phi_4(\zeta) = \left( \frac{2m_1}{\varepsilon \sqrt{\Delta} \cos(2\sqrt{-m_1} \zeta) - m_2} \right)^{\frac{1}{2}}. \quad (2.9)$$

When  $m_1 > 0, \Delta < 0$ ,

$$\Phi_5(\zeta) = \left( \frac{2m_1}{\varepsilon \sqrt{-\Delta} \sinh(2\sqrt{m_1} \zeta) - m_2} \right)^{\frac{1}{2}}. \quad (2.10)$$

When  $m_1 < 0, \Delta > 0$ ,

$$\Phi_6(\zeta) = \left( \frac{2m_1}{\varepsilon \sqrt{\Delta} \sin(2\sqrt{-m_1} \zeta) - m_2} \right)^{\frac{1}{2}}. \quad (2.11)$$

When  $m_1 > 0, m_3 > 0$ ,

$$\Phi_7(\zeta) = \left( \frac{-m_1 \operatorname{sech}^2(\sqrt{m_1} \zeta)}{m_2^2 - 2\varepsilon \sqrt{m_1 m_3} \tanh(\sqrt{m_1} \zeta)} \right)^{\frac{1}{2}}. \quad (2.12)$$

When  $m_1 < 0, m_3 > 0$ ,

$$\Phi_8(\zeta) = \left( \frac{-m_1 \sec^2(\sqrt{-m_1} \zeta)}{m_2^2 + 2\varepsilon \sqrt{-m_1 m_3} \tan(\sqrt{-m_1} \zeta)} \right)^{\frac{1}{2}}. \quad (2.13)$$

When  $m_1 > 0, m_3 > 0$ ,

$$\Phi_9(\zeta) = \left( \frac{m_1 \operatorname{csch}^2(\sqrt{m_1} \zeta)}{m_2^2 + 2\varepsilon \sqrt{m_1 m_3} \coth(\sqrt{m_1} \zeta)} \right)^{\frac{1}{2}}. \quad (2.14)$$

When  $m_1 < 0, m_3 > 0$ ,

$$\Phi_{10}(\zeta) = \left( \frac{-m_1 \csc(\sqrt{-m_1} \zeta)}{m_2^2 + 2\varepsilon \sqrt{-m_1 m_3} \tanh(\sqrt{-m_1} \zeta)} \right)^{\frac{1}{2}}, \quad (2.15)$$

When  $m_1 > 0, \Delta = 0$ ,

$$\Phi_{11}(\zeta) = \left( -\frac{m_1}{m_2} (1 + \varepsilon \tanh(\frac{\sqrt{m_1}}{2} \zeta)) \right)^{\frac{1}{2}}. \quad (2.16)$$

When  $m_1 > 0, \Delta = 0$ ,

$$\Phi_{12}(\zeta) = \left( -\frac{m_1}{m_2} (1 + \varepsilon \coth(\frac{\sqrt{m_1}}{2} \zeta)) \right)^{\frac{1}{2}}. \quad (2.17)$$

When  $m_1 > 0$ ,

$$\Phi_{13}(\zeta) = 4 \left( \frac{m_1 e^{2\varepsilon\sqrt{m_1}\zeta}}{(e^{2\varepsilon\sqrt{m_1}\zeta} - 4m_2)^2 - 64m_1m_3} \right)^{\frac{1}{2}}. \quad (2.18)$$

When  $m_1 > 0, m_2 = 0$ ,

$$\Phi_{14}(\zeta) = 4 \left( \frac{\pm m_1 e^{2\varepsilon\sqrt{m_1}\zeta}}{(1 - 64m_1m_3 e^{4\varepsilon\sqrt{m_1}\zeta})} \right)^{\frac{1}{2}}, \quad (2.19)$$

where  $\Delta = m_2^2 - 4m_1m_3$  and  $\varepsilon = \pm 1$ .

**Step 3.** Finally, the ansatz (2.4) is substituted into (2.3) and all the coefficients of all powers of  $\Phi(\zeta)$  are equated to zero. The resultant set of algebraic equations are investigated to find the relations among the parameters particularly for unknowns  $m_1, m_2, m_3, n_i (i = 1, 2, 3, \dots, N)$ .

### 3. Application:

Consider the complex-valued wave transformation

$$w(x, t) = W(\xi) \exp(i\psi(x, t)), \quad (3.1)$$

with  $\xi = x - vt$ ,  $\psi(x, t) = -kx + \omega t + \theta$ ,  $W(\zeta)$  representing various features of the wave and the phase  $\psi(x, t)$  of the soliton, the other constants  $k, \theta, \omega, v$  are frequency, the number of wave, the constant of wave and soliton speed, respectively. Putting (3.1) into (2.2), followed by an uncoupled system of equations derived from the real and imaginary parts of the equation.

The imaginary component reads:

$$v = -\alpha - 2a_1k + c_1W^2 - (2\mu m + (2m+1)\lambda)W^{2m}, \quad (3.2)$$

which anticipates the soliton speed. By setting  $m=1$  in Eq. (3.2) and equating the coefficients of linearly independent functions to zero forces:

$$v = \alpha - 2a_1k, c_1 = 2\mu + 3\lambda, \quad (3.3)$$

The real component reads:

$$aW'' - (\alpha k + W + a_1k^2)W - k\lambda W^{2m+1} - c_1kW^3 + b_1W^5 = 0, \quad (3.4)$$

which furnishes soliton profile.

Moreover Eq. (3.4) is reduced to following ODE, by employing dependent variable change as  $W(\xi) = \sqrt{U(\xi)}$ ,

$$\begin{aligned} -4U^2(\alpha k + w + a_1k^2) + 4(-c_1kU^3 + b_1U^4) - 4k\lambda U^{2m+1} \\ + a_1(-(U')^2 + 2UU'') = 0, \end{aligned} \quad (3.5)$$

Taking  $m=1$ , by homogeneous balancing method we consider  $U''$  or  $U'^2$  with  $U^4$  in equation (3.5), we got  $N=1$  and proceeded as

$$U(\xi) = n_0 + n_1 \Phi(\xi) \tag{3.6}$$

With  $n_0$  and  $n_1$  being constant terms to be determine. Substituting Eq. (3.6) into Eq. (3.5) by the fact that Eq. (2.5) is satisfied and equating the coefficients of  $\Phi^i(\xi)$  for  $(i=0,1,2,\dots,N)$  to zero we get the algebraic system of equations as follows:

$$\begin{aligned} const: & -4n_0^2\alpha k - 4n_0^2ak^2 - 4ckn_0^3 - 4\lambda kn_0^3 - 4n_0^3\varpi + 4bn_0^4 = 0, \\ \Phi(\xi): & -8n_0n_1ak^2 - 8n_0n_1\varpi - 8n_0n_1\alpha k + 16bn_0^3n_1 - 12\lambda kn_0^2n_1 \\ & + 2an_1n_0m_1 - 12ckn_0^2n_1 = 0, \\ \Phi^2(\xi): & -12\lambda kn_0n_1^2 + 24bn_0^2n_1^2 - 4n_1^2\varpi - 4n_1^2\alpha k - 4n_1^2ak^2 \\ & + an_1^2m_1 - 12ckn_0n_1^2 = 0, \\ \Phi^3(\xi): & -4ckn_1^3 - 4\lambda kn_1^3 + 16bn_0n_1^3 + 4an_1n_0m_2 = 0, \\ \Phi^4(\xi): & 4bn_1^4 + 3an_1^2m_2 = 0, \\ \Phi^5(\xi): & 6an_1n_0m_3 = 0, \\ \Phi^6(\xi): & 5an_1^2m_3 = 0. \end{aligned}$$

Solving this set of algebraic equations using *Maple*, we found our solutions to be

$$\begin{aligned} n_0 &= \frac{3}{8} \frac{k(c+\lambda)}{b}, \quad n_1 = n_1, \quad m_1 = \frac{3}{16} \frac{k^2(c+\lambda)^2}{ab}, \quad m_2 = -\frac{4}{3} \frac{bn_1^2}{a}, \\ m_3 &= 0, \quad \varpi = -\frac{1}{64} \frac{k(30\lambda ck + 15\lambda^2k + 15c^2k + 64bak + 64ab)}{b}. \end{aligned}$$

When  $ab > 0$ ,

$$\begin{aligned} w_1(x,t) &= \sqrt{\frac{3k(\lambda+c)}{8b} \left( 1 + \operatorname{sech} \left( \frac{k(\lambda+c)}{4} \sqrt{\frac{3}{ab}} (x-vt) \right) \right)} \\ &\times \exp(i(-kx - \frac{1}{64} \frac{k(30\lambda ck + 15\lambda^2k + 15c^2k + 64bak + 64ab)}{b} t + \theta)), \\ w_2(x,t) &= \sqrt{\frac{3k(c+\lambda)}{8b} \left( 1 + i \operatorname{csch} \left( \frac{k(c+\lambda)}{4} \sqrt{\frac{3}{ab}} (x-vt) \right) \right)} \\ &\times \exp(i(-kx - \frac{1}{64} \frac{k(30\lambda ck + 15\lambda^2k + 15c^2k + 64bak + 64ab)}{b} t + \theta)), \end{aligned}$$

$$\begin{aligned}
w_3(x,t) &= \sqrt{\frac{3k(c+\lambda)}{8b} \left( 1 + \sqrt{\frac{2}{1 + \cosh\left(\frac{k(c+\lambda)}{2} \sqrt{\frac{3}{ab}}(x-vt)\right)}} \right)} \\
&\times \exp\left(i\left(-kx - \frac{1}{64} \frac{k(30\lambda ck + 15\lambda^2 k + 15c^2 k + 64bak + 64\alpha b)}{b} t + \theta\right)\right), \\
w_4(x,t) &= \sqrt{\frac{3k(\lambda+c)}{8b} \left( 1 + \sqrt{\frac{2}{i \sinh\left(\frac{k(\lambda+c)}{2} \sqrt{\frac{3}{ab}}(x-vt)\right) + 1}} \right)} \\
&\times \exp\left(i\left(-kx - \frac{1}{64} \frac{k(30\lambda ck + 15\lambda^2 k + 15c^2 k + 64bak + 64\alpha b)}{b} t + \theta\right)\right), \\
w_5(x,t) &= \sqrt{\frac{3k(c+\lambda)}{8b} + n_1 k(c+\lambda)} \sqrt{\frac{3e^{\frac{k(c+\lambda)}{2} \sqrt{\frac{3}{ab}}(x-vt)}}}{ab \left( e^{\frac{k(c+\lambda)}{2} \sqrt{\frac{3}{ab}}(x-vt)} + \frac{16bn_1^2}{3a} \right)^2}} \\
&\times \exp\left(i\left(-kx - \frac{1}{64} \frac{k(30\lambda ck + 15\lambda^2 k + 15c^2 k + 64bak + 64\alpha b)}{b} t + \theta\right)\right),
\end{aligned}$$

When  $ab < 0$ ,

$$\begin{aligned}
w_6(x,t) &= \sqrt{\frac{3k(c+\lambda)}{8b} \left( 1 + \sqrt{\frac{2}{1 + \cos\left(\frac{k(c+\lambda)}{2} \sqrt{\frac{3}{ab}}(x-vt)\right)}} \right)} \\
&\times \exp\left(i\left(-kx - \frac{1}{64} \frac{k(30\lambda ck + 15\lambda^2 k + 15c^2 k + 64bak + 64\alpha b)}{b} t + \theta\right)\right), \\
w_7(x,t) &= \sqrt{\frac{3k(\lambda+c)}{8b} \left( 1 + \sqrt{\frac{2}{\left(\sin\left(\frac{k(\lambda+c)}{2} \sqrt{\frac{3}{ab}}(x-vt)\right) + 1\right)}} \right)} \\
&\times \exp\left(i\left(-kx - \frac{1}{64} \frac{k(30\lambda ck + 15\lambda^2 k + 15c^2 k + 64bak + 64\alpha b)}{b} t + \theta\right)\right).
\end{aligned}$$


#### 4. Physical Illustrations

In this section, we illustrate some particular forms of the solutions determined above for particular choices of parameters satisfying given constraints.


The real and imaginary components of the solution  $w_1(x,t)$  is depicted in Fig 1a-1b for some particular choice of the parameters as  $k = 1, c = -1, \lambda = -1, a = 1, b = 1, v = 1, \alpha = 1, \theta = 0$  in the finite domain of independent variables  $x \in [-10,10] \times t \in [0,2]$ . Fig 1c represent the modulus of the same solution for the same parameters and in the same domain. Both the real imaginary components represent some periodic solutions as the modulus is a single solitary wave propagating to the right along the space axis.

In Fig 2, we plot the solution  $w_2(x,t)$  in the finite domain  $x \in [-10,10] \times t \in [0,2]$  for the parameters  $k = 1, c = -1, \lambda = -1, a = 1, b = 1, v = 1, \alpha = 1, \theta = 0$ . The real and imaginary components of the solutions have some positive and negative explosions at some points of space and time due to their functional nature, Fig 2a-b. The modulus of  $w_2(x,t)$  also indicates explosions with huge positive amplitudes time to time owing to the behaviours of the real and imaginary components of the solutions.


Fig 3 demonstrates the solution  $w_6(x,t)$  for the particular parameters  $k = 1, c = -1/2, \lambda = -1, a = -1, b = 1, v = 1, \alpha = 1, \theta = 0$ . In Fig 3a-b, the real and imaginary components represent multi waves having positive or negative jumps time to time. The modulus of this solution represent periodic long waves with positive amplitudes, Fig 3c.



(a) Real component




(b) Imaginary component




(c) Modulus


Fig 1. The solution  $w_1(x,t)$  for  $k = 1, c = -1, \lambda = -1, a = 1, b = 1, v = 1, \alpha = 1, \theta = 0$



(a) Real component




(b) Imaginary component




(c) Modulus

Fig 2. The solution  $w_2(x, t)$  for  $k = 1, c = -1, \lambda = -1, a = 1, b = 1, v = 1, \alpha = 1, \theta = 0$




(a) Real component



(b) Imaginary component





(c) Modulus

Fig 3. The solution  $w_6(x, t)$  for  $k = 1, c = -1/2, \lambda = -1, a = -1, b = 1, \nu = 1, \alpha = 1, \theta = 0$

## 5. Conclusion:

In this paper, we found exact travelling wave solutions for the non-linear perturbed GIE with the help of new EAEM. The complex valued solutions represent traveling waves in various forms. Even though some are of the form of well-known bell shaped multi waves, the shape of some others are completely different from them.

Eventually, the scheme is simple, direct, efficient and robust for solving perturbed Gerdjikov–Ivanov non-linear equation and it kind that usually arises experimental and mathematical physics.

## References

- [1] Biswas, A., Rezazadeh, H., Mirzazadeh, M., Eslami, M., Zhou, Q., Moshokoa, S. P., & Belic, M. (2018). Optical solitons having weak non-local non-linearity by two integration schemes. *Optik*, 164, 380-384.
- [2] Mirzazadeh, M., Eslami, M., & Biswas, A. (2014). Soliton solutions of the generalized Klein–Gordon equation by using (G'/G)-expansion method. *Computational and Applied Mathematics*, 33(3), 831-839.
- [3] Rezazadeh, H. (2018). New solitons solutions of the complex Ginzburg-Landau equation with Kerr law non-linearity. *Optik*, 167, 218-227.
- [4] Rezazadeh, H., Mirhosseini-Alizamini, S. M., Eslami, M., Rezazadeh, M., Mirzazadeh, M., & Abbagari, S. (2018). New optical solitons of non-linear conformable fractional Schrödinger-Hirota equation. *Optik*, 172, 545-553.
- [5] Eslami, M., & Rezazadeh, H. (2016). The first integral method for Wu–Zhang system with conformable time-fractional derivative. *Calcolo*, 53(3), 475-485.
- [6] Eslami, M., & Mirzazadeh, M. (2014). First integral method to look for exact solutions of a variety of Boussinesq-like equations. *Ocean Engineering*, 83, 133-137.
- [7] Eslami, M., & Mirzazadeh, M. (2016). Optical solitons with Biswas–Milovic equation for power law and dual-power law non-linearities. *Non-linear Dynamics*, 83(1-2), 731-738.
- [8] Eslami, M. (2016). Trial solution technique to chiral non-linear Schrodinger's equation in (1+2)-dimensions. *Non-linear Dynamics*, 85(2), 813-816.

- [9] Rezazadeh, H., Korkmaz, A., Eslami, M., Vahidi, J., & Asghari, R. (2018). Traveling wave solution of conformable fractional generalized reaction Duffing model by generalized projective Riccati equation method. *Optical and Quantum Electronics*, 50(3), 150.
- [10] Khodadad, F. S., Nazari, F., Eslami, M., & Rezazadeh, H. (2017). Soliton solutions of the conformable fractional Zakharov-Kuznetsov equation with dual-power law non-linearity. *Optical and Quantum Electronics*, 49(11), 384.
- [11] Zhou, Q., Ekici, M., Sonmezoglu, A., Mirzazadeh, M., & Eslami, M. (2016). Optical solitons with Biswas–Milovic equation by extended trial equation method. *Non-linear Dynamics*, 84(4), 1883-1900.
- [12] Jamilu S., Abdullahi J., & Abubakar M. G. (2018). New exact solution for the (3+1) conformable space-time fractional modified Korteweg–de-Vries equations via Sine-Cosine Method, *Journal of Taibah University for Science*, DOI:10.1080/16583655.2018.1537642,1-5.
- [13] Lakhveer Kaura, Abdul-Majid Wazwaz, (2018). Optical solitons for perturbed GIE, *Optik - International Journal for Light and Electron Optics* 174, 447–451.
- [14] Osman, M. S., Korkmaz, A., Rezazadeh, H., Mirzazadeh, M., Eslami, M., & Zhou, Q. (2018). The Unified Method for Conformable Time Fractional Schrödinger Equation with Perturbation Terms. *Chinese Journal of Physics*, 56(5), 2500-2506.
- [15] Korkmaz, A. (2017). Exact solutions to (3+ 1) conformable time fractional Jimbo–Miwa, Zakharov–Kuznetsov and modified Zakharov–Kuznetsov equations. *Communications in Theoretical Physics*, 67(5), 479.
- [16] Korkmaz, A. (2018). On the wave solutions of conformable fractional evolution equations. *Communications*, 67, 68-79.
- [17] Korkmaz, A., & Hosseini, K. (2017). Exact solutions of a non-linear conformable time-fractional parabolic equation with exponential non-linearity using reliable methods. *Optical and Quantum Electronics*, 49(8), 278.
- [18] Korkmaz, A. (2018). Complex Wave Solutions to Mathematical Biology Models I: Newell–Whitehead–Segel and Zeldovich Equations. *Journal of Computational and Non-linear Dynamics*, 13(8), 081004.
- [19] Fan, E. (2000). Integrable evolution systems based on GIEs, bi-Hamiltonian structure, finite-dimensional integrable systems and N-fold Darboux transformation. *Journal of Mathematical Physics*, 41(11), 7769-7782.
- [20] Dai, H. H., & Fan, E. G. (2004). Variable separation and algebro-geometric solutions of the GIE. *Chaos, Solitons & Fractals*, 22(1), 93-101.
- [21] Manafian, J., & Lakestani, M. (2016). Optical soliton solutions for the Gerdjikov–Ivanov model via  $\tan(\phi/2)$ -expansion method. *Optik-International Journal for Light and Electron Optics*, 127(20), 9603-9620.
- [22] Guo, L., Zhang, Y., Xu, S., Wu, Z., & He, J. (2014). The higher order rogue wave solutions of the GIE. *Physica Scripta*, 89(3), 035501.
- [23] Kakei, S., & Kikuchi, T. (2005). Solutions of a derivative non-linear Schrödinger hierarchy and its similarity reduction. *Glasgow Mathematical Journal*, 47(A), 99-107.
- [24] Xu, S., & He, J. (2012). The rogue wave and breather solution of the Gerdjikov-Ivanov equation. *Journal of Mathematical Physics*, 53(6), 063507.
- [25] Sirendaoreji. (2006). A new auxiliary equation and exact travelling wave solutions of non-linear equations. *Physics Letters A*, 356(2), 124-130.